

On Yukawa quasi-unification with $\mu < 0$

M.E. Gómez,^{1,*} G. Lazarides,^{2,†} and C. Pallis^{3,‡}

¹*Centro de Física das Interações Fundamentais (CFIF), Departamento de Física,
Instituto Superior Técnico, Av. Rovisco Pais, Lisboa 1049-001, Portugal*

²*Physics Division, School of Technology, Aristotle University of Thessaloniki, Thessaloniki 54124, Greece*

³*Scuola Internazionale Superiore Di Studi Avanzati (SISSA), Via Beirut 2-4, Trieste 34013, Italy*

(Dated: February 1, 2008)

Although recent data on the muon anomalous magnetic moment strongly disfavor the constrained minimal supersymmetric standard model with $\mu < 0$, they cannot exclude it because of theoretical ambiguities. We consider this model supplemented by a Yukawa quasi-unification condition which allows an acceptable b -quark mass. We find that the cosmological upper bound on the lightest sparticle relic abundance is incompatible with the data on the branching ratio of $b \rightarrow s\gamma$, which is evaluated by including all the next-to-leading order corrections. Thus, this scheme is not viable.

PACS numbers: 12.10.Kt, 12.60.Jv, 95.35.+d

The constrained minimal supersymmetric standard model (CMSSM), which is a predictive version of the minimal supersymmetric standard model (MSSM) based on universal boundary conditions, can be further restricted by being embedded in a supersymmetric (SUSY) grand unified theory (GUT) with a gauge group containing $SU(4)_c$ and $SU(2)_R$. This can lead [1] to Yukawa unification [2], i.e., the exact unification of the third generation ‘asymptotic’ Yukawa couplings. However, for both signs of the MSSM parameter μ , the CMSSM supplemented by the assumption of exact Yukawa unification yields unacceptable values of the b -quark mass. This is due to the generation of sizeable SUSY corrections to m_b [3] which have the sign of μ . The predicted tree-level $m_b(M_Z)$, which turns out to be close to the upper edge of its 95% confidence level (c.l.) experimental range $2.684 - 3.092$ GeV, receives, for $\mu > 0$ ($\mu < 0$), large positive (negative) corrections which drive it well above (a little below) the allowed range. This range is derived from the 95% c.l. range for $m_b(m_b)$ in the $\overline{MS}$ scheme $3.95 - 4.55$ GeV [4] evolved up to M_Z in accord with the analysis of Ref. [5] and with $\alpha_s(M_Z) = 0.1185$.

In Ref. [6], we have built SUSY GUT models based on the Pati-Salam gauge group $SU(4)_c \times SU(2)_L \times SU(2)_R$ which moderately violate Yukawa unification and, thus, can allow an acceptable b -quark mass for both signs of μ even with universal boundary conditions. These models yield a class of ‘asymptotic’ Yukawa quasi-unification conditions (YQUCs) replacing exact Yukawa unification. Supplementing the CMSSM with one of these YQUCs, we achieved [6], for $\mu > 0$, an acceptable $m_b(M_Z)$ in accord with all the phenomenological and cosmological requirements. The corresponding SUSY GUT model leads [7] to a new version of shifted hybrid inflation [8].

Here, we will examine whether the CMSSM with a

YQUC can yield an acceptable $m_b(M_Z)$ in accord with all the available constraints for $\mu < 0$ too. In this case, compatibility of the b -quark mass with the data requires a deviation from Yukawa unification which is much smaller than the one needed for $\mu > 0$. Consequently, there is no need to extend the simplest SUSY GUT model of Ref. [6], which yields a suppressed violation of Yukawa unification originating from non-renormalizable interactions.

The CMSSM with $\mu < 0$ is now strongly disfavored (but not excluded [9]) by the constraint on the deviation, δa_μ , of the measured value of the muon anomalous magnetic moment a_μ from its predicted value in the standard model (SM) a_μ^{SM} . The recent data [10] on a_μ confirm the earlier ones [11] but their precision is twice that of the previous data. On the other hand, a_μ^{SM} is not yet stabilized. This is mainly due to the hadronic vacuum polarization contribution. According to the most updated [12] evaluation of this contribution, there is a considerable discrepancy between the results based on e^+e^- data and on τ data. Taking into account both results, we obtain the 95% c.l. range $-4.7 \times 10^{-10} \lesssim \delta a_\mu \lesssim 56 \times 10^{-10}$, where the negative values are allowed only by the τ -based calculation. Using the formulae of Ref. [13], we calculate δa_μ in the CMSSM and we observe that the result is negative (positive) for $\mu < 0$ ($\mu > 0$) and increases (decreases) as the lightest sparticle (LSP) mass m_{LSP} increases. Consequently, a lower bound on m_{LSP} can be derived from the lower (upper) experimental limit on δa_μ for $\mu < 0$ ($\mu > 0$). Also, we see that $\mu < 0$ is allowed only by the τ -based calculation.

The CMSSM with $\mu < 0$ is, also, severely restricted by the recent experimental results [14] on the inclusive branching ratio $\text{BR}(b \rightarrow s\gamma)$. We compute this branching ratio by using an updated version of the relevant calculation in the micrOMEGAs code [15]. Comparing carefully this version with our calculation of $\text{BR}(b \rightarrow s\gamma)$ in Ref. [6], we concluded that this improved version is complete and reliable. The SM contribution is calculated by using the formalism of Ref. [16] and incorporating the improvements of Ref. [17]. The charged Higgs boson ($H^\pm$) contribution is evaluated by including the next-to-

*Electronic address: mgomez@cfif.ist.utl.pt

†Electronic address: lazaride@eng.auth.gr

‡Electronic address: pallis@sisssa.it

leading order (NLO) QCD corrections from Ref. [18] and the $\tan\beta$ enhanced contributions from Ref. [19]. The dominant SUSY contribution includes resummed NLO SUSY QCD corrections from Ref. [19], which hold for large $\tan\beta$. The $H^\pm$ contribution interferes constructively with the SM contribution, while the SUSY one interferes constructively (destructively) with the other two contributions when $\mu < 0$ ($\mu > 0$). The SM plus $H^\pm$ and the SUSY contributions decrease as m_{LSP} increases and so, for $\mu < 0$, an additional lower bound on m_{LSP} can be derived [20] from the upper experimental limit on this branching ratio. At 95% c.l., the experimental range of $\text{BR}(b \rightarrow s\gamma)$ is estimated [6] by combining appropriately the various experimental and theoretical errors involved. The result is $1.9 \times 10^{-4} \lesssim \text{BR}(b \rightarrow s\gamma) \lesssim 4.6 \times 10^{-4}$. We find that the NLO corrections reduce $\text{BR}(b \rightarrow s\gamma)$ by 10 – 15%. In Ref. [21] (which adopts the opposite sign convention for μ), where the $\tan\beta$ enhanced and NLO SUSY QCD corrections were ignored, the reduction was considerably higher yielding a less stringent restriction.

The LSP mass is bounded above by the requirement that the LSP relic abundance $\Omega_{\text{LSP}} h^2$ in the universe does not exceed the upper limit on the cold dark matter (CDM) abundance derived from observations. Recent results [22] suggest that, at 95% c.l., $\Omega_{\text{LSP}} h^2 \lesssim 0.22$. In general, $\Omega_{\text{LSP}} h^2$ increases with m_{LSP} . So, its value is expected to be quite large due to the large m_{LSP} 's required by the constraints from δa_μ and, especially, $b \rightarrow s\gamma$. The $\Omega_{\text{LSP}} h^2$ is calculated by using `micrOMEGAs` [15], which is the most complete code available at present. It includes accurately thermally averaged exact tree-level cross sections of all possible (co)annihilation processes and treats pole effects properly with one-loop QCD corrected decay widths. In the model which we will consider, the LSP is an almost pure bino ($\tilde{\chi}$) and the next-to-LSP (NLSP) is the lightest stau ($\tilde{\tau}_2$). Important annihilation channels are not only the ones with fermions $f\bar{f}$ in the final state, but also the ones with HZ , $W^\pm H^\mp$, hA [23] (h , H are the CP-even neutral Higgs bosons and A the CP-odd neutral Higgs boson). On the other hand, as we can observe from the sparticle spectrum of the model (see Fig. 1), m_A is much smaller than $2m_{\text{LSP}}$. So, the LSP annihilation via the s-channel exchange of an A -boson is not enhanced as in the $\mu > 0$ case [6, 24]. As a consequence, the only available mechanism for reducing $\Omega_{\text{LSP}} h^2$ is bino-stau coannihilation [6, 21, 25, 26] (for an updated analysis, see Ref. [27]), which is activated when the relative mass splitting between the $\tilde{\tau}_2$ and the LSP, $\Delta_{\tilde{\tau}_2} = (m_{\tilde{\tau}_2} - m_{\text{LSP}})/m_{\text{LSP}}$, is reduced below about 0.25. For fixed m_{LSP} , $\Omega_{\text{LSP}} h^2$ decreases with $\Delta_{\tilde{\tau}_2}$, since coannihilation becomes more efficient. So the CDM bound yields an upper limit on $\Delta_{\tilde{\tau}_2}$.

We will consider here the CMSSM with $\mu < 0$ which results from the simplest SUSY GUT model constructed in Ref. [6] without the extra Higgs superfields $\bar{\phi}$, ϕ , $\bar{\phi}'$, ϕ' . To reduce the number of free parameters, we will further assume that the non-renormalizable $SU(2)_R$ triplet coupling of Ref. [6] which leads to violation of Yukawa




FIG. 1: The mass parameters m_A , m_0 , $M_{1/2}$ and M_{SUSY} versus m_{LSP} for $A_0 = 0$, $\Delta_{\tilde{\tau}_2} = 0$ and $m_b(M_Z) = 2.888$ GeV.

unification is suppressed. The violation is then achieved predominantly via the non-renormalizable $SU(2)_R$ singlet coupling of Ref. [6] and, thus, $\alpha_1 = \alpha_2$ in Eq. (15) of this reference, with α_1 , α_2 of order 0.1 or smaller. Note that the other alternative with the triplet coupling dominating, which was studied in Ref. [6] with $\mu > 0$, is not viable for $\mu < 0$ since it leads to a charged LSP. Under the above assumptions, the YQUC takes the form:

$$h_t : h_b : h_\tau = |1 - c| : |1 - c| : |1 + 3c|, \quad (1)$$

where c is a complex parameter of order 0.1 or smaller. We will take $c < 0$, where the maximal splitting between h_b , h_τ and, thus, the largest enhancement of $m_b(M_Z)$ is achieved for fixed $|c| \neq 0$. For $-1/3 < c < 0$, this splitting is $-\delta h \equiv (h_b - h_\tau)/h_b = -4c/(1 - c)$. We see that the tau Yukawa coupling becomes smaller than the top and bottom Yukawa couplings which remain unified, in contrast to the $\mu > 0$ case of Ref. [6], where the tau and bottom Yukawa couplings split from the top Yukawa coupling by the same amount. The SUSY GUT model used leads [8] to successful shifted hybrid inflation.

Below the GUT scale M_{GUT} , this SUSY GUT model reduces to the CMSSM with universal soft SUSY breaking scalar masses m_0 , gaugino masses $M_{1/2}$, and trilinear scalar couplings A_0 at M_{GUT} . It also implies the YQUC in Eq. (1). We closely follow the renormalization group and radiative electroweak breaking analysis of Ref. [6]. The conditions for electroweak breaking are imposed on an optimized scale $M_{\text{SUSY}} = (m_{\tilde{t}_1} m_{\tilde{t}_2})^{1/2}$ ($\tilde{t}_{1,2}$ are the stop mass eigenstates), where the sparticle spectrum is also evaluated. We incorporate the two-loop corrections to the CP-even neutral Higgs boson mass matrix by using `FeynHiggsFast` [28] and the SUSY corrections to $m_b(M_{\text{SUSY}})$ and $m_\tau(M_{\text{SUSY}})$ from Ref. [29]. The corrections to $m_\tau(M_{\text{SUSY}})$ lead [21] to a small enhancement of $\tan\beta$. For $m_t(m_t) = 166$ GeV, $m_\tau(M_Z) = 1.746$ GeV, and any given $m_b(M_Z)$ in its 95% c.l. range (2.684 – 3.092 GeV), we can determine the parameters c and $\tan\beta$ at M_{SUSY} so that the YQUC in




FIG. 2: Restrictions on the $m_{\text{LSP}} - \Delta\tilde{\tau}_2$ plane for $A_0 = 0$ and $2.684 \text{ GeV} \lesssim m_b(M_Z) \lesssim 3.092 \text{ GeV}$. The solid (dot-dashed) line is the lower bound on m_{LSP} from $\delta a_\mu \gtrsim -4.7 \times 10^{-10}$ ($\text{BR}(b \rightarrow s\gamma) \lesssim 4.6 \times 10^{-4}$). The dashed (double dot-dashed) line is the upper bound on m_{LSP} ($\Delta\tilde{\tau}_2$) from $\Omega_{\text{LSP}} h^2 \lesssim 0.22$.

Eq. (1) is satisfied. We are then left with only three free input parameters m_0 , $M_{1/2}$ and A_0 . The first two can be replaced [21] by m_{LSP} and $\Delta\tilde{\tau}_2$. In Fig. 1, we present the values of m_A , m_0 , $M_{1/2}$ and M_{SUSY} versus m_{LSP} for $A_0 = 0$, $\Delta\tilde{\tau}_2 = 0$, $m_b(M_Z) = 2.888 \text{ GeV}$. These values are affected very little by varying $m_b(M_Z)$. Our scheme, unlike the $\mu > 0$ model of Ref. [6], yields $2m_{\text{LSP}} \gg m_A$.

The restrictions on the $m_{\text{LSP}} - \Delta\tilde{\tau}_2$ plane for $A_0 = 0$ and any $m_b(M_Z)$ in its 95% c.l. range are given in Fig. 2. The lower bound on m_{LSP} from $\delta a_\mu \gtrsim -4.7 \times 10^{-10}$ corresponds to $m_b(M_Z) \simeq 3.092 \text{ GeV}$ and is represented by a solid line. The maximal $\Delta\tilde{\tau}_2$ on this line is about 0.0105 and yields $\Omega_{\text{LSP}} h^2 \simeq 0.22$ for the same value of $m_b(M_Z)$. As $m_b(M_Z)$ decreases, the maximal $\Delta\tilde{\tau}_2$ from $\Omega_{\text{LSP}} h^2 \lesssim 0.22$ increases along the double dot-dashed line and reaches its overall maximal value $\Delta\tilde{\tau}_2 \simeq 0.0142$ at $m_b(M_Z) \simeq 2.684 \text{ GeV}$. The upper bound on m_{LSP} from $\Omega_{\text{LSP}} h^2 \lesssim 0.22$ is achieved at $m_b(M_Z) \simeq 2.684 \text{ GeV}$ and corresponds to the dashed line. In the lightly shaded region allowed by δa_μ and CDM considerations, $612.4 \text{ GeV} \lesssim m_{\text{LSP}} \lesssim 873.4 \text{ GeV}$. The maximal m_{LSP} is achieved at $\Delta\tilde{\tau}_2 = 0$ yielding $\text{BR}(b \rightarrow s\gamma) \simeq 5.26 \times 10^{-4}$. The lower bound on m_{LSP} (dot-dashed line) from $\text{BR}(b \rightarrow s\gamma) \lesssim 4.6 \times 10^{-4}$ corresponds to $m_b(M_Z) \simeq 3.092 \text{ GeV}$. In the corresponding allowed (dark shaded) area, $m_{\text{LSP}} \gtrsim 1305.04 \text{ GeV}$ with the minimal m_{LSP} achieved at $\Delta\tilde{\tau}_2 = 0$ and yielding $\Omega_{\text{LSP}} h^2 \simeq 0.65$. So, for $A_0 = 0$, there is no region where all the restrictions are satisfied. Note that the constraints $\text{BR}(b \rightarrow s\gamma) \gtrsim 1.9 \times 10^{-4}$ and $\delta a_\mu \lesssim 56 \times 10^{-10}$ always hold for the CMSSM with $\mu < 0$. Also, $m_h \gtrsim 114.4 \text{ GeV}$ [30] is valid due to the heavy SUSY spectrum.

Departure from the $A_0 = 0$ case does not change our conclusion as we can see from Fig. 3, where the restrictions on the $m_{\text{LSP}} - A_0/M_{1/2}$ plane are presented for any $\Delta\tilde{\tau}_2$ and any allowed $m_b(M_Z)$. Actually, we can just




FIG. 3: Restrictions on the $m_{\text{LSP}} - A_0/M_{1/2}$ plane for any $\Delta\tilde{\tau}_2$ and $2.684 \text{ GeV} \lesssim m_b(M_Z) \lesssim 3.092 \text{ GeV}$. The notation is generally as in Fig. 2. The dotted line and the inclined parts of the solid line are explained in the main text.

take $\Delta\tilde{\tau}_2 = 0$, where all the constraints happen to become as less restrictive as possible for any given A_0 (see Fig. 2). We follow the same notation for the various lines and areas as in Fig. 2. For $-1.84 \lesssim A_0/M_{1/2} \lesssim 1.21$ ($-1.67 \lesssim A_0/M_{1/2} \lesssim 1.03$), the solid (dashed) line corresponds to $\delta a_\mu \simeq -4.7 \times 10^{-10}$ ($\Omega_{\text{LSP}} h^2 \simeq 0.22$) with $m_b(M_Z) \simeq 3.092 \text{ GeV}$ (2.684 GeV). As $|A_0|$ increases, m_A decreases and reaches its lowest allowed value ($\simeq 100 \text{ GeV}$ for the $\tan\beta$'s encountered here [31]) on the boundaries of the lightly shaded regions for fixed A_0 . Allowing $m_b(M_Z)$ to increase from its lower value, we then obtain the upper and lower inclined parts of the dashed line corresponding to $m_A \simeq 100 \text{ GeV}$. Their end points lie at $A_0/M_{1/2} \simeq 1.21, -1.84$, where $m_b(M_Z)$ is maximized. At the maximal (minimal) m_{LSP} on the dotted line (inclined parts of the solid line), $\Omega_{\text{LSP}} h^2 \lesssim 0.22$ ($\delta a_\mu \gtrsim -4.7 \times 10^{-10}$) is not saturated, due to the combined requirements $m_A \gtrsim 100 \text{ GeV}$ and $m_{\tilde{\tau}_2} \gtrsim m_{\tilde{\chi}}$.

The maximal m_{LSP} in the lightly shaded area of Fig. 3, which is allowed by δa_μ and CDM considerations, and for $A_0/M_{1/2} > 0$ (< 0) is $m_{\text{LSP}} \simeq 868 \text{ GeV}$ (967.3 GeV) and is achieved at $A_0/M_{1/2} \simeq 1.03$ (-1.67). This corresponds to the upper (lower) corner of the dashed line, where $\text{BR}(b \rightarrow s\gamma) \simeq 5.58 \times 10^{-4}$ (5.45×10^{-4}). Note that, for $A_0/M_{1/2} < 0$, the processes $\tilde{\tau}_2 \tilde{\tau}_2^* \rightarrow W^\pm H^\mp, ZA$ become more efficient. Their total contribution to the effective cross section is $3 - 6.5\%$ as $A_0/M_{1/2}$ decreases from 0 to -1.67 . So, coannihilation is strengthened and bigger m_{LSP} 's are allowed than for $A_0/M_{1/2} > 0$. The minimal m_{LSP} in the dark shaded area, allowed by $b \rightarrow s\gamma$, is $m_{\text{LSP}} \simeq 1295 \text{ GeV}$ and is achieved at $A_0/M_{1/2} \simeq -0.2$, where $\Omega_{\text{LSP}} h^2 \simeq 0.62$. Thus, even for $A_0 \neq 0$, there is no range of parameters allowed by all the constraints.

In the lightly shaded area of Fig. 3, $\tan\beta$ (c) ranges between about 44.1 (-0.09) and 51.5 (-0.007). These values are achieved at the lowest corner of this area ($A_0/M_{1/2} \simeq -1.89$, $m_b(M_Z) \simeq 3.092 \text{ GeV}$) and the

upper corner of the dashed line ($A_0/M_{1/2} \simeq 1.03$, $m_b(M_Z) \simeq 2.684$ GeV) respectively. Consequently, the splitting $-\delta h$ ranges from about 0.33 to 0.028. However, fixing $m_b(M_Z)$ to its central value, the range of $-\delta h$ is reduced to $0.13-0.09$ corresponding to $47.6 \lesssim \tan \beta \lesssim 49$ and $-0.038 \lesssim c \lesssim -0.024$.

The present investigation is an improved version of our analysis in Ref. [21] (the sign of μ there is opposite to the one adopted here). The main improvements are the replacement of Yukawa unification by the YQUC of Eq. (1) in connection with the more stringent present bounds on m_b , the consideration of the $\delta\alpha_\mu$ constraint [13], the inclusion of the NLO SUSY QCD and $\tan\beta$ enhanced corrections [19] in $\text{BR}(b \rightarrow s\gamma)$, and the evaluation of $\Omega_{\text{LSP}}h^2$ by the updated code of Ref. [15]. These improvements turn out to be of crucial importance overruling our previous conclusion. Our results are essentially unaffected by allowing $\alpha_s(M_Z)$ to vary in its 95% c.l. range, which slightly enlarges [5, 6] the range of $m_b(M_Z)$. This is due to the fact that, here, the LSP annihilation to $b\bar{b}$

via an A -pole exchange is subdominant in $\Omega_{\text{LSP}}h^2$.

In summary, we studied the CMSSM with $\mu < 0$ and a YQUC from the simplest SUSY GUT of Ref. [6]. We imposed the constraints from m_b , $\delta\alpha_\mu$, $b \rightarrow s\gamma$ and CDM. Although the NLO corrections to $b \rightarrow s\gamma$ (coannihilations) drastically reduce (enhance) the lower (upper) bound on m_{LSP} , the $b \rightarrow s\gamma$ and CDM requirements remain incompatible. Thus, despite the fact that, with the τ -based calculation of α_μ^{SM} , the $\delta\alpha_\mu$ and CDM criteria can be simultaneously valid, this model is excluded.

We are grateful to the micrOMEGAs team, G. Bélanger, F. Boudjema, A. Pukhov and A. Semenov, for providing us with their updated code for calculating $\text{BR}(b \rightarrow s\gamma)$. We also thank J. Ellis, W. Liao, A. Masiero and E. Palante for useful discussions. This work was supported by European Union under the RTN contracts HPRN-CT-2000-00148 and HPRN-CT-2000-00152. M.E.G. acknowledges support from the ‘Fundação para a Ciência e Tecnologia’ under contract SFRH/BPD/5711/2001.

-
- [1] G. Lazarides and C. Panagiotakopoulos, Phys. Lett. B **337**, 90 (1994); S. Khalil, G. Lazarides and C. Pallis, *ibid.* **508**, 327 (2001).
 - [2] B. Ananthanarayan, G. Lazarides and Q. Shafi, Phys. Rev. D **44**, 1613 (1991); Phys. Lett. B **300**, 245 (1993).
 - [3] M. Carena, M. Olechowski, S. Pokorski and C.E.M. Wagner, Nucl. Phys. B **426**, 269 (1994); R. Hempfling, Phys. Rev. D **49**, 6168 (1994); L. Hall, R. Rattazzi and U. Sarid, *ibid.* **50**, 7048 (1994).
 - [4] C.T. Sachrajda, Nucl. Instrum. Meth. A **462**, 23 (2001).
 - [5] H. Baer, J. Ferrandis, K. Melnikov and X. Tata, Phys. Rev. D **66**, 074007 (2002).
 - [6] M.E. Gómez, G. Lazarides and C. Pallis, Nucl. Phys. B **638**, 165 (2002).
 - [7] R. Jeannerot, S. Khalil and G. Lazarides, J. High Energy Phys. **07**, 069 (2002).
 - [8] R. Jeannerot, S. Khalil, G. Lazarides and Q. Shafi, J. High Energy Phys. **10**, 012 (2000); G. Lazarides, in *Recent Developments in Particle Physics and Cosmology*, edited by G.C. Branco, Q. Shafi and J.I. Silva-Marcos (Kluwer Acad. Pub., Dordrecht, 2001) p. 399 (hep-ph/0011130); R. Jeannerot, S. Khalil and G. Lazarides, in *The Proceedings of Cairo International Conference on High Energy Physics*, edited by S. Khalil, Q. Shafi and H. Tallaat (Rinton Press, Princeton, 2001) p. 254 (hep-ph/0106035).
 - [9] S.P. Martin and J.D. Wells, hep-ph/0209309.
 - [10] G.W. Bennett *et al.*, (Muon (g-2) Collaboration), Phys. Rev. Lett. **89**, 101804 (2002); **89**, 129903(E) (2002).
 - [11] H.N. Brown *et al.*, (Muon (g-2) Collaboration), Phys. Rev. Lett. **86**, 2227 (2001).
 - [12] M. Davier, S. Eidelman, A. Höcker and Z. Zhang, hep-ph/0208177.
 - [13] S.P. Martin and J.D. Wells, Phys. Rev. D **64**, 035003 (2001).
 - [14] R. Barate *et al.* (ALEPH Collaboration), Phys. Lett. B **429**, 169 (1998); K. Abe *et al.* (BELLE Collaboration), *ibid.* **511**, 151 (2001); S. Chen *et al.* (CLEO Collaboration), Phys. Rev. Lett. **87**, 251807 (2001).
 - [15] G. Bélanger, F. Boudjema, A. Pukhov and A. Semenov, Comput. Phys. Commun. **149**, 103 (2002).
 - [16] A.L. Kagan and M. Neubert, Eur. Phys. J. C **7**, 5 (1999).
 - [17] P. Gambino and M. Misiak, Nucl. Phys. B **611**, 338 (2001).
 - [18] M. Ciuchini, G. Degrossi, P. Gambino and G. Giudice, Nucl. Phys. B **527**, 21 (1998).
 - [19] G. Degrossi, P. Gambino and G.F. Giudice, J. High Energy Phys. **12**, 009 (2000).
 - [20] F. M. Borzumati, M. Olechowski and S. Pokorski, Phys. Lett. B **349**, 311 (1995); H. Baer, M. Brhlik, D. Castaño and X. Tata, Phys. Rev. D **58**, 015007 (1998).
 - [21] M.E. Gómez, G. Lazarides and C. Pallis, Phys. Lett. B **487**, 313 (2000).
 - [22] C. Pryke *et al.*, Astrophys. J. **568**, 46 (2002).
 - [23] T. Nihei, L. Roszkowski and R. Ruiz de Austri, J. High Energy Phys. **05**, 063 (2001).
 - [24] J. Ellis, T. Falk, G. Gani, K.A. Olive and M. Srednicki, Phys. Lett. B **510**, 236 (2001).
 - [25] J. Ellis, T. Falk and K.A. Olive, Phys. Lett. B **444**, 367 (1998); J. Ellis, T. Falk, K.A. Olive and M. Srednicki, Astropart. Phys. **13**, 181 (2000); **15**, 413(E) (2001).
 - [26] M.E. Gómez, G. Lazarides and C. Pallis, Phys. Rev. D **61**, 123512 (2000).
 - [27] T. Nihei, L. Roszkowski and R. Ruiz de Austri, J. High Energy Phys. **07**, 024 (2002).
 - [28] S. Heinemeyer, W. Hollik and G. Weiglein, hep-ph/0002213.
 - [29] D. Pierce, J. Bagger, K. Matchev and R. Zhang, Nucl. Phys. B **491**, 3 (1997); M. Carena, D. Garcia, U. Nierste and C.E.M. Wagner, *ibid.* **577**, 88 (2000); S.F. King and M. Oliveira, Phys. Rev. D **63**, 015010 (2001).
 - [30] ALEPH, DELPHI, L3 and OPAL Collaborations, The LEP working group for Higgs boson searches, hep-ex/0107029.
 - [31] J.A. Valls (CDF Collaboration), FERMILAB-CONF-99/263-E.